

Canonical BCS Approximation for the Attractive Hubbard Model

K. Tanaka and F. Marsiglio

Department of Physics, University of Alberta, Edmonton, Alberta, Canada T6G 2J1

Abstract. We test the canonical BCS wave functions for fixed number of electrons for the attractive Hubbard model. We present results in one dimension for various chain lengths, electron densities, and coupling strengths. The ground-state energy and energy gap to the first excited state are compared with the exact solutions obtained by the Bethe ansatz as well as with the results from the conventional grand-canonical BCS approximations. While the canonical and grand-canonical BCS results are both in very good agreement with the exact results for the ground state energies, improvements due to conserving the electron number in finite systems are manifest in the energy gap. As the system size is increased, the canonical results converge to the grand-canonical ones. The “parity” effect that arises from the number parity (even or odd) of electrons are studied with our canonical scheme.

In recent experiments on nanoscale superconducting Al particles (e.g., Ref. [1]), the number of electrons in a particle was fixed by charging effects and the excitation spectrum has been measured by tunneling of a single electron. They have observed the number “parity” effect, the difference in the excitation spectra that arises due to the number of electrons in a particle being even or odd. For these systems, the validity of the BCS theory that utilizes the grand canonical ensemble is questionable. We address this issue by using the attractive Hubbard model, which serves as the minimal model to describe s-wave superconductivity. We test the canonical BCS wave function that has a fixed number of electrons for this system. While our formulation can be applied for larger systems up to three dimensions, we present results for one dimension, for various system sizes, electron densities and coupling strengths. Exact solutions are available in one dimension via Bethe Ansatz techniques [2], and the grand canonical BCS solutions have recently been evaluated for large system sizes [3].

The attractive Hubbard Hamiltonian is given by

$$H = - \sum_{\substack{i,\delta \\ \sigma}} t_{\delta} (a_{i+\delta,\sigma}^{\dagger} a_{i\sigma} + \text{h.c.}) - |U| \sum_i n_{i\uparrow} n_{i\downarrow} \quad (1a)$$

$$= \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}} a_{\mathbf{k}\sigma}^{\dagger} a_{\mathbf{k}\sigma} - \frac{|U|}{N} \sum_{\mathbf{k}\mathbf{k}'} a_{\mathbf{k}\uparrow}^{\dagger} a_{-\mathbf{k}+\mathbf{l}\downarrow}^{\dagger} a_{-\mathbf{k}'+\mathbf{l}\downarrow} a_{\mathbf{k}'\uparrow} , \quad (1b)$$

where $a_{i\sigma}^\dagger$ ($a_{i\sigma}$) creates (annihilates) an electron with spin σ at site i and $n_{i\sigma}$ is the number operator, $n_{i\sigma} = a_{i\sigma}^\dagger a_{i\sigma}$ (i is the index for the primitive vector $\mathbf{R}_i$). The t_δ is the hopping rate of electrons from one site to a neighbouring site $\mathbf{R}_\delta$ away. In this work, we include the nearest neighbours only, and $t \equiv t_\delta$. The $|U|$ is the coupling strength between electrons on the same site, and we have explicitly included the fact that the interaction is attractive. In Eq.(1b), we have Fourier transformed the Hamiltonian with periodic boundary conditions for N sites in each dimension. The kinetic energy is given by

$$\epsilon_{\mathbf{k}} = -2 \sum_{\delta} t_{\delta} \cos \mathbf{k} \cdot \mathbf{R}_{\delta} , \quad (2)$$

where $\mathbf{R}_{\delta}$ is the coordinate vector connecting sites i to $i + \delta$.

We perform variational calculations using the component of the BCS wave function that has a given number of pairs ν and thus conserves the number of electrons $N_e = 2\nu$. The BCS wave function is a superposition of pair states with all the possible numbers of pairs $\{\nu\}$:

$$|\text{BCS}\rangle_{\text{GC}} = \prod_{\mathbf{k}} (u_{\mathbf{k}} + v_{\mathbf{k}} a_{\mathbf{k}\uparrow}^\dagger a_{-\mathbf{k}\downarrow}^\dagger) |0\rangle \equiv \sum_{\nu=0}^{\infty} |\Psi_{\nu}\rangle , \quad (3)$$

where $|0\rangle$ denotes the vacuum state, and $|\Psi_0\rangle \equiv |0\rangle$. The ν -pair component $|\Psi_{\nu}\rangle$ can be obtained by rearranging $|\text{BCS}\rangle_{\text{GC}}$ into a power series of the pair creation operator $a_{\mathbf{k}\uparrow}^\dagger a_{-\mathbf{k}\downarrow}^\dagger$ and can be written as

$$|\Psi_{2\nu}\rangle = \frac{1}{\nu!} \prod_{i=1}^{\nu} \left(\sum_{\mathbf{k}_i} g_{\mathbf{k}_i} a_{\mathbf{k}_i\uparrow}^\dagger a_{-\mathbf{k}_i\downarrow}^\dagger \right) |0\rangle , \quad (4)$$

where $\mathbf{k}_i \neq \mathbf{k}_j$ for all $i \neq j$, and we have defined $g_{\mathbf{k}_i} = \left(\prod_{\mathbf{k}} u_{\mathbf{k}} \right)^{1/\nu} v_{\mathbf{k}_i} / u_{\mathbf{k}_i}$. An alternative way of formulating the problem is to express the above wave function as a particle-number projection of $|\text{BCS}\rangle_{\text{GC}}$ [4–6]. The wave function with an odd number of electrons $N_e = 2\nu + 1$ is defined by

$$|\Psi_{2\nu+1}\rangle = a_{\mathbf{q}\sigma}^\dagger \frac{1}{\nu!} \prod_{i=1}^{\nu} \left(\sum_{\mathbf{k}_i} g_{\mathbf{k}_i} a_{\mathbf{k}_i\uparrow}^\dagger a_{-\mathbf{k}_i\downarrow}^\dagger \right) |0\rangle , \quad (5)$$

with $\mathbf{k}_i \neq \mathbf{k}_j$ for all $i \neq j$. We construct the energy gap by the definition

$$\Delta_{N_e} = \frac{1}{2} (E_{N_e-1} - 2E_{N_e} + E_{N_e+1}) . \quad (6)$$

In Fig. 1 we show the ground state energy as a function of the electron density $n = N_e/N$ for $N = 16$, for $|U|/t = 10$ and 4. The improvements of the canonical over the grand canonical energies can be seen, while the former with even N_e 's converge to the latter for smaller n and larger $|U|$, and both results are in very good agreement




FIGURE 1. Ground state energy as a function of the electron density for $N = 16$.




FIGURE 2. Energy gap as a function of the electron density for $N = 32$ and $|U|/t = 2$.

with the exact ones for small n . The difference between the energies with even and odd N_e 's is apparent for large $|U|$.

The energy gap (scaled by the kinetic energy parameter t) is shown in Fig. 2 as a function of the electron density, for $N = 32$ and $|U|/t = 2$. The canonical method improves the grand canonical one significantly for small system size and weak coupling. As the system size or the coupling strength is increased, the canonical results converge towards the grand canonical ones. For $|U|/t = 2$, $N = 32$ is large enough that the canonical gaps are closer to the grand canonical ones than to the exact solutions, and the canonical and grand canonical gaps are more or less converged for small n . The gap for odd N_e is negative. Its magnitude is different from the even N_e gap for larger density and weak coupling. Also in this limit, the even gap oscillates as a function of the density. Both of these effects result from the quantized (unperturbed) energy levels due to finite size, and become negligible for strong coupling and for large system size (small density).




FIGURE 3. Occupation probability for different k states as a function of the coupling strength for $N = 64$ and $\nu = 32$.

Finally, in Fig. 3, we show the occupation probability of different $\mathbf{k}$ states as a function of the coupling strength. In the strong coupling limit, all the unperturbed states are more or less equally occupied (in Fig. 3, this limit is not reached yet). As the coupling strength goes to zero, the distribution function for free electrons is recovered. Though the figure shown has been produced by the canonical scheme, the grand canonical approximation also yields the same probability distribution for such a large system size.

In conclusion, the canonical method yields improvements in the energy gap for small system size and weak coupling. Further study is being made for comparison with the grand canonical scheme that conserves the number parity [7].

Acknowledgments

F. M. acknowledges support from NSERC of Canada, and the Canadian Institute for Advanced Research. K. T. acknowledges generous support from the Avadh Bhatia Fellowship. We thank Bob Teshima for suggestions to improve the numerical efficiency for evaluating required integrals. Calculations were performed on the 42 node SGI parallel processor at the University of Alberta.

REFERENCES

1. Black, C.T., Ralph, D.C., and Tinkham, M., *Phys. Rev. Lett.* **76**, 688 (1996).
2. Lieb, E.H., and Wu, F.Y., *Phys. Rev. Lett.* **20**, 1445 (1968).
3. Marsiglio, F., *Phys. Rev. B* **55**, 575 (1997).
4. Dietrich, K., Mang, H.J., and Pradal, J.H., *Phys. Rev.* **135**, B22 (1964).
5. Braun, F., and von Delft, J., *Phys. Rev. Lett.* **81**, 4712 (1998).
6. Tanaka, K., and Marsiglio, F., cond-mat/9902173
7. Jankó, B., Smith, A., and Ambegaokar, V., *Phys. Rev. B* **50**, 1152 (1994).